


# CHAPTER – 8 BASIC CONCEPTS OF DIFFERENTIAL AND INTEGRAL CALCULUS

## (A) DIFFERENTIAL CALCULUS



<b>INTRODUCTION</b>	Differentiation is one of the most important fundamental operations in calculus. Its theory primarily depends on the idea of limit and continuity of function.
<b>DERIVATIVE OR DIFFERENTIAL COEFFICIENT</b>	Let $y = f(x)$ be a function. If $h$ be the small increment in $x$ and the corresponding increment in $y$ or $f(x)$ be $y = f(x+h) - f(x)$
<b>STANDARD FORMULAS</b>	

$\frac{d}{dx}(a) = 0$	$\frac{d}{dx}[\ln x] = \frac{d}{dx}[\log_e x] = \frac{1}{x} \frac{dx}{dx}$
$\frac{d}{dx}(x) = 1$	$\frac{d}{dx}[\log_a x] = \log_a e \frac{1}{x} \frac{dx}{dx}$
$\frac{d}{dx}(au) = a \frac{du}{dx}$	$\frac{d}{dx} e^u = e^u \frac{du}{dx}$
$\frac{d}{dx}(u + v - w) = \frac{du}{dx} + \frac{dv}{dx} - \frac{dw}{dx}$	$\frac{d}{dx} a^u = a^u \ln a \frac{du}{dx}$
$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$	$\frac{d}{dx}(u^v) = v u^{v-1} \frac{du}{dx} + \ln u \cdot u^v \frac{dv}{dx}$
$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{1}{v} \frac{du}{dx} - \frac{u}{v^2} \frac{dv}{dx}$	$\frac{d}{dx} \sin u = \cos u \frac{du}{dx}$
$\frac{d}{dx}(u^n) = n u^{n-1} \frac{du}{dx}$	$\frac{d}{dx} \cos u = -\sin u \frac{du}{dx}$
$\frac{d}{dx}(\sqrt{u}) = \frac{1}{2\sqrt{u}} \frac{du}{dx}$	$\frac{d}{dx} \tan u = \sec^2 u \frac{du}{dx}$
$\frac{d}{dx}\left(\frac{1}{u}\right) = -\frac{1}{u^2} \frac{du}{dx}$	$\frac{d}{dx} \cot u = -\csc^2 u \frac{du}{dx}$
$\frac{d}{dx}\left(\frac{1}{u^n}\right) = -\frac{n}{u^{n+1}} \frac{du}{dx}$	$\frac{d}{dx} \sec u = \sec u \tan u \frac{du}{dx}$
$\frac{d}{dx}[f(u)] = \frac{d}{du}[f(u)] \frac{du}{dx}$	$\frac{d}{dx} \csc u = -\csc u \cot u \frac{du}{dx}$

**IMPLICIT FUNCTIONS**

A function in the form  $f(x, y) = 0$ . For example,  $x^2y^2 + 3xy + y = 0$  where  $y$  cannot be directly defined as a function of  $x$  is called an implicit function of  $x$ .


**PARAMETRIC EQUATION**

When both the variables  $x$  and  $y$  are expressed in terms of a parameter (a third variable), the involved equations are called parametric equations. For the parametric equations  $x = f(t)$  and  $y = h(t)$  the differential coefficient  $\frac{dy}{dx}$

**LOGARITHMIC DIFFERENTIATION**

The process of finding out derivative by taking logarithm in the first instance is called logarithmic differentiation.

**GEOMETRIC INTERPRETATION OF THE DERIVATIVE**



**COST FUNCTION**

Total cost consists of two parts (i) Variable Cost (ii) Fixed Cost.

	<b>Average cost (AC or C )</b>	$\frac{\text{Total Cost } C(X)}{\text{Output } \bar{X}}$
	<b>Average variable cost (AVC)</b>	$\frac{\text{Variable Cost } V(X)}{\text{Output } \bar{X}}$
	<b>Average Fixed Cost (AFC)</b>	$\frac{\text{Fixed Cost } F(X)}{\text{Output } \bar{X}}$
<b>MARGINAL COST</b>	If $C(x)$ the total cost producing $x$ units then the increase in cost in producing one more unit is called marginal cost at an output level of $x$ units	
<b>REVENUE FUNCTION</b>	Revenue, $R(x)$ , gives the total money obtained (Total turnover) by selling units of a product. If $x$ units are sold at $P$ per unit, then $R(x) = P \cdot X$	
<b>PROFIT FUNCTION</b>	Profit $P(x)$ , the difference of between total revenue $R(x)$ and total Cost $C(x)$ .	

**(B) INTEGRAL CALCULUS**

<b>INTEGRATION</b>	<pre> graph TD     f_prime["f'(x)"] --&gt; I["INTEGRATION"]     f_prime --&gt; D["DIFFERENTIATION"]     I --&gt; f["f(x)"]     D --&gt; f             </pre>	<p>Integration is the reverse process of differentiation.</p>
--------------------	--	---

## DEFINITE INTEGRATION

**DEFINITION** The collection of all antiderivatives of  $f$  is called the **indefinite integral** of  $f$  with respect to  $x$ , and is denoted by

$$\int f(x) dx.$$

The symbol  $\int$  is an **integral sign**. The function  $f$  is the **integrand** of the integral, and  $x$  is the **variable of integration**.

### Differentiation Formulas:

1.  $\frac{d}{dx}(x) = 1$
2.  $\frac{d}{dx}(ax) = a$
3.  $\frac{d}{dx}(x^n) = nx^{n-1}$
4.  $\frac{d}{dx}(\cos x) = -\sin x$
5.  $\frac{d}{dx}(\sin x) = \cos x$
6.  $\frac{d}{dx}(\tan x) = \sec^2 x$
7.  $\frac{d}{dx}(\cot x) = -\csc^2 x$
8.  $\frac{d}{dx}(\sec x) = \sec x \tan x$
9.  $\frac{d}{dx}(\csc x) = -\csc x(\cot x)$
10.  $\frac{d}{dx}(\ln x) = \frac{1}{x}$
11.  $\frac{d}{dx}(e^x) = e^x$
12.  $\frac{d}{dx}(a^x) = (\ln a)a^x$
13.  $\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$
14.  $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$
15.  $\frac{d}{dx}(\sec^{-1} x) = \frac{1}{|x|\sqrt{x^2-1}}$

### Integration Formulas:

1.  $\int 1 dx = x + C$
2.  $\int a dx = ax + C$
3.  $\int x^n dx = \frac{x^{n+1}}{n+1} + C, n \neq -1$
4.  $\int \sin x dx = -\cos x + C$
5.  $\int \cos x dx = \sin x + C$
6.  $\int \sec^2 x dx = \tan x + C$
7.  $\int \csc^2 x dx = -\cot x + C$
8.  $\int \sec x(\tan x) dx = \sec x + C$
9.  $\int \csc x(\cot x) dx = -\csc x + C$
10.  $\int \frac{1}{x} dx = \ln |x| + C$
11.  $\int e^x dx = e^x + C$
12.  $\int a^x dx = \frac{a^x}{\ln a} + C, a > 0, a \neq 1$
13.  $\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + C$
14.  $\int \frac{1}{1+x^2} dx = \tan^{-1} x + C$
15.  $\int \frac{1}{|x|\sqrt{x^2-1}} dx = \sec^{-1} x + C$

### Properties of Definite Integral

Assuming  $f$  and  $g$  are continuous functions

$$\int_a^b f(x)dx = \int_b^a f(x)dx$$

$$\int_a^a f(x)dx = 0$$

$$\int_a^b c dx = c(b - a), \text{ where } c \text{ is any constant}$$

$$\int_a^b cf(x)dx = c \int_a^b f(x)dx, \text{ where } c \text{ is any constant}$$

$$\int_a^b [f(x) + g(x)]dx = \int_a^b f(x)dx + \int_a^b g(x)dx$$

$$\int_a^b [f(x) - g(x)]dx = \int_a^b f(x)dx - \int_a^b g(x)dx$$

$$\int_a^b f(x)dx + \int_b^c f(x)dx = \int_a^c f(x)dx$$



#### Question 1

Find an expression for  $y$  given  $\frac{dy}{dx} = 7x^5$

(a) 6

(b) 2

(c) 3

(d) 5

**Answer: a**

**Explanation:**

$$\frac{dy}{dx} = 7x^5 \rightarrow dy = 7x^5 dx$$

Integrating both sides, we have

$$\int dy = \int 7x^5 dx \rightarrow y = \frac{7x^6}{6} + c$$

#### Question 2

Find an expression for  $y$  given  $\frac{dy}{dx} = x^{-\frac{3}{4}}$

(a)  $\frac{2}{3}$

(b)  $\frac{1}{4}$

(c)  $\frac{5}{4}$

(d) None

**Answer: b****Explanation:**

$$\frac{dy}{dx} = x^{-3/4}$$

$$Y = \frac{x^{-3/4+1}}{-\frac{3}{4}+1} = \frac{x^{1/4}}{1/4}$$

$$Y = 4x^{1/4}$$

**Question 3** **$-12x^{-4} = \int -12x^{-4} - dx$  solve it;**

(a) 6

(b) 2

(c) 3

(d) 4

**Answer: d****Explanation:**

$$dy = \int -12x^{-4} dx$$

$$= -12 \int x^{-4} dx$$

$$\text{Use } \int ax^n dx = a \int x^n dx = \frac{ax^{n+1}}{n+1} + c$$

$$= + \left( \frac{-12x^{-3}}{-3} \right) + c$$

$$n = -4, n + 1 = -4 + 1 = -3$$

$$Y = 4x^{-3} + c$$

$$\text{Simplifying fraction, } \frac{-12}{3} = 4$$

**Question 4****Given  $f'(x) = \frac{1}{2}x^{\frac{1}{3}} - \frac{1}{4}x^{\frac{1}{4}} + \pi$ , find  $f(x)$** 

(a) 6

(b) 2

(c) 3

(d) None

**Answer: d****Explanation:**

$$\frac{1}{2}x^{\frac{1}{3}} - \frac{1}{4}x^{\frac{1}{4}} + \pi$$

$$\int \frac{1}{2}x^{\frac{1}{3}} dx - \int \frac{1}{4}x^{\frac{1}{4}} dx + \int \pi dx$$

$$\begin{aligned} & \frac{1}{2} \int x^{\frac{1}{3}} dx - \frac{1}{4} \int x^{\frac{1}{4}} dx + \pi \int dx \\ &= \frac{\frac{1}{2} x^{\frac{4}{3}}}{\frac{4}{3}} - \frac{\frac{1}{4} x^{\frac{5}{4}}}{\frac{5}{4}} + \frac{\pi x}{2} + C \\ &= \frac{3x^{\frac{4}{3}}}{8} - \frac{1}{4} \times \frac{4}{5} x^{\frac{5}{4}} + \pi x + C \end{aligned}$$

**Question 5**

Given  $f'(x) = \int \left( \frac{2}{x} + \frac{3}{x^2} + \frac{1}{x^5} \right) dx$

(a) -6

(b) 2

(c) -4

(d) None

**Answer: c****Explanation:**

$$\begin{aligned} & \int \left\{ \frac{2}{x} + \frac{3}{x^2} + \frac{1}{x^5} \right\} dx \\ &= \int \left( \frac{2}{x} + 3x^{-2} + x^{-5} \right) dx \quad \text{write as negative exponence} \\ &= \int \frac{2}{x} dx + \int 3x^{-2} dx + \int x^{-5} dx \quad \text{Use } \int f(x) dx + g(x) dx = \int f(x) dx + \int g(x) dx \\ &= 2 \ln |x| + \frac{3x^{-1}}{-1} + \frac{x^{-4}}{-4} + c \quad \text{Use } \int ax^n dx = a \int x^n dx = \frac{ax^{n+1}}{n+1} + c \\ &= 2 \ln |x| - \frac{3}{x} - \frac{1}{4x^4} + c \quad \text{Simplify } \frac{3}{-1} \end{aligned}$$

**Question 6**

Integrate  $\int \frac{3}{x^{\frac{1}{2}}} dx$

(a)  $6\sqrt{x+c}$ (b)  $\sqrt{x+c}$ (c)  $8\sqrt{x+c}$ (d)  $9\sqrt{x+c}$ **Answer: a****Explanation:**

$$\begin{aligned}\int \frac{3}{x^{\frac{1}{2}}} dx &= \int 3x^{-1/2} \\ &= \frac{3x^{-1/2+1}}{-\frac{1}{2}+1} + C \\ &= \frac{3x^{\frac{1}{2}}}{\frac{1}{2}} + C \\ &= 6x^{\frac{1}{2}} + C \\ &= 6\sqrt{x} + C\end{aligned}$$

**Question 7**

Find  $y$  as a function of  $x$  if  $\frac{d^2y}{dx^2} = 2x$  when  $x = 2, y = 7$

(a)  $y = \frac{x^3}{3} + c$

(b)  $y = \frac{x^2}{3} + c$

(c)  $y = \frac{x}{3} + c$

(d) None

**Answer: a****Explanation:**

$$\begin{aligned}\int 2x dx &= 2 \int x dx \\ &= \left( \frac{2x^{1+1}}{1+1} \right) + C\end{aligned}$$

$$\text{Use } \int ax^n dx = a \int x^n dx = \frac{ax^{n+1}}{n+1} + C$$

$$\frac{dy}{dx} = x^2 + c$$

Multiply of fraction/simplify

$$\text{Finding } y = \int \frac{dy}{dx} = \int x^2 dx$$

$$Y = \frac{x^3}{3} + c$$

$$\text{Use } \int x^n dx = \frac{1}{n+1} x^{n+1} + c$$

At (2, 7)

$$7 = \frac{2^3}{3} + c$$

Substituting  $x = 2$  and  $y = 7$  to find  $c$ 

$$C = \frac{21}{8}$$

Thus, the function is  $y = \frac{x^3}{3} + c$ .

**Question 8**

Integrate  $\int \left( w + \frac{1}{w} \right) \left( w - \frac{1}{w} \right) dx$

(a)  $\frac{w^3}{3} + \frac{1}{w}$

(b)  $\frac{w^3}{3} + \frac{1}{w} + c$



(c)  $\frac{w}{3} + \frac{1}{w} + c$

(d) None

**Answer: b****Explanation:**

$$\int \left(w + \frac{1}{w}\right) \left(w - \frac{1}{w}\right) dw$$

$$= \int \left(w^2 - \frac{1}{w^2}\right) dw$$

squares

$$= \int w^2 dw - \int \frac{1}{w^2} dw$$

$$\int g(x) dx$$

$$= \int w^2 dw - \int w^{-2} dw$$

$$= \frac{w^3}{3} + \frac{1}{w} + c$$

Express the product as a difference of two

Use  $\int f(x) dx + g(x) dx = \int f(x) dx +$

Express in negative exponential form

Use  $\int x^n dx = \frac{1}{n+1} x^{n+1} + c$ . Simplify

**Question 9**If  $\frac{d^2y}{dx^2} = 10 - 3x$ , find  $\frac{dy}{dx} + c$ 

(a)  $10x - \frac{3}{2}x^2 + c$

(b)  $10x - \frac{3}{2} + c$

(c)  $10 - \frac{3}{2}x^2 + c$

(d) none

**Answer: a****Explanation:**

$$\frac{dy}{dx} = \int (10 - 3x) dx = \int 10 dx - \int 3x dx$$

Use  $\int f(x) dx + g(x) dx =$

$$\int f(x) dx + \int g(x) dx$$

$$= 10x - \left(\frac{3x^{1+1}}{1+1}\right) + c$$

Use  $\int ax^n dx = a \int x^n dx = \frac{ax^{n+1}}{n+1} + c$

$$= 10x - \left(\frac{3x^2}{2}\right) + c$$

Simplify

$$= 10x - \frac{3}{2}x^2 + c$$

**Question 10**Calculate  $\int x^7 dx$ 

(a)  $\frac{1}{8}x^7 + c$

(b)  $\frac{1}{7}x^7 + c$

(c)  $\frac{1}{8}x^8 + c$

(d) None

**Answer: c**

**Explanation:**

$$\int x^7 dx = \frac{1}{7+1} x^{7+1} + c \quad \text{Use } \int x^n dx = \frac{1}{n+1} x^{n+1} + c \text{ and substitute } n = 7$$

$$= \frac{1}{8} x^8 + c$$

**Question 11**

If  $\int f(x) dx = xe^{-\log|x|} + f(x)$ , then  $f(x)$  is

- (a) 1 (b) 0  
(c)  $ce^x$  (d)  $\log x$

**Answer: c****Explanation:**

$$\int f(x) dx = xe^{\log\left|\frac{1}{x}\right|} + f(x) \Rightarrow \int f(x) dx = \frac{x}{|x|} + f(x)$$

On differentiating both sides, we get

$$F(x) = 0 + f'(x) \quad \text{we know}$$

$$\frac{d}{dx}(e^x) = e^x, \therefore f(x) = ce^x$$

**Question 12**

If  $f(t) = \int_{-t}^t \frac{dx}{1+x^2}$ , then  $f'(1)$  is

- (a) 0 (b)  $\frac{2}{3}$   
(c) -1 (d) 1

**Answer: d****Explanation:**

$$\text{Given } f(t) = \int_{-t}^t \frac{dx}{1+x^2} = [\tan^{-1}x]_{-t}^t = 2 \tan^{-1}t$$

$$\text{Differentiating with respect to } f'(t) = \frac{2}{1+t^2}$$

$$\therefore f'(1) = \frac{2}{2} = 1$$

**Question 13**

The existence of first order partial derivatives implies continuity

(a) True

(b) False

(c) Not sure

(d) Invalid Question

**Answer: b****Explanation:**

The mere existence cannot be declared as a condition for continuity because the second order derivatives should also be continuous.

**Question 14**

$y = (x^2(1 + x^3))$  find  $\frac{dy}{dx}$

(a)  $-(2x + 5x^4)\sin(x^2 + x^2)$ (b)  $(2x + 5x^4)\sin(x^2 + x^5)$ (c)  $(2x + 5x^4)(x^2 + x^5)$ 

(d) none

**Answer: d****Explanation:**

$\frac{dy}{dx} = -\sin(x^2 + x^5) \frac{d}{dx}(x^2 + x^5)$  using the chain rule

$\frac{dy}{dx} = -(\sin(x^2 + x^5))(2x + 5x^4)$  using the basic derivatives

$\frac{dy}{dx} = -(2x + 5x^4)\sin(x^2 + x^5)$  reordering factors

**Question 15**

If  $f(x) = x^k$  and  $f'(1) = 10$ , then the value of k is

(a) 10

(b) -10

(c)  $\frac{1}{10}$ 

(d) None

**Answer: a****Explanation:**

$F(x) = x^k$

$F(1) = f(1) = k \times 1$

$10 = k \times 1$

$K = 10$

**Question 16**

The points of discontinuity of the function,  $f(x) = \frac{x^2+2x+5}{x^2-3x+2}$  are

- (a)  $x=0, x=1$  (b)  $x=1, x=2$   
 (c)  $x=0, x=2$  (d) None

**Answer: b****Explanation:**

$$f(x) = \frac{x^2+2x+5}{x^2-3x+2}$$

$$\text{Denominator} = 0$$

$$X^2 - 3x + 2 = 0$$

$$(x-1)(x-2) = 0$$

$$X = 1, x = 2$$

**Question 17**

The gradient of a function is parallel to the velocity vector of the level curve

- (a) True (b) False  
 (c) Not sure (d) Invalid questions

**Answer: b****Explanation:**

The gradient is perpendicular and not parallel to the velocity vector of the level curve.

**Question 18**

$$y = (8 + x^3)(x^3 - 8)$$

- (a)  $6x^5$  (b)  $x^5$   
 (c)  $6x$  (d) None

**Answer: a****Explanation:**

This problem is solvable as a product but if you realize that you are looking at a difference of two squares, it becomes very simple.

$$Y = (8 + x^3)(x^3 - 8) = x^6 - 64$$

$$\frac{dy}{dx} = 6x^5$$

**Question 19**

If  $(x, y, z, t) = xy + zt + x^2 y z t$ ;  $x = k^3$ ;  $y = k^2$ ;  $z = k$ ;  $t = \sqrt{k}$

Find  $\frac{df}{dt}$  at  $k = 1$

- (a) 34 (b) 16  
(c) 32 (d) 61

**Answer: b**

**Explanation:**

Using chain rule we have

$$\begin{aligned} \frac{df}{dt} &= f_x \frac{dx}{dk} + f_y \frac{dy}{dk} + f_z \frac{dz}{dk} + f_t \frac{dt}{dk} \\ &= (y + 2xyzt).(3k^2) + (x + x^2zt).(2k) + (t + x^2yt).(1) + (z + x^2yz).\left(\frac{1}{2\sqrt{k}}\right) \end{aligned}$$

Put  $k = 1$ ; we have  $x=y=z=t=1$

$$9 + 4 + 2 + 1 = 16.$$

**Question 20**

If  $(x, y) = x^2 + y^3$ ;  $x = t^2 + t^3$ ;  $y = t^3 + t^9$  find  $\frac{df}{dt}$  at  $t=1$ .

- (a) 0 (b) 1  
(c) -1 (d) 164

**Answer: d**

**Explanation:**

Using chain rule we have

$$\begin{aligned} \frac{df}{dt} &= f_x \frac{dx}{dt} + f_y \frac{dy}{dt} \\ &= (2x).(2t + 3t^2) + (3y^2).(3t^2 + 9t^8) \end{aligned}$$

Put  $t = 1$ ; we have  $x = 2$ ;  $y = 2$

$$= 4.(5) + 12.(12) = 164.$$

**Question 21**

$f(x, y) = x^2 + xyz + z$  find  $f_x$  at  $(1, 1, 1)$

- (a) 0 (b) 1  
(c) 3 (d) -1

**Answer: c**

**Explanation:**

$$F_x = 2x + yz$$

$$\text{Put } (x, y, z) = (1, 1, 1)$$

$$F_x = 2 + 1 = 3.$$

**Question 22**

**Necessary condition of Euler's theorem is**

**(a) z should be homogenous and of order n**      **(b) x should not homogeneous but for order n**

**(c) Should be implicit**      **(d) should be the function of x and y only**

**Answer: a**

**Explanation:**

Of x and y of order 'n' then  $x \frac{dz}{dx} + y \frac{dz}{dy} = nz$

Answer 'b' is incorrect as z should be homogeneous.

Answer 'c' is incorrect as z should not be implicit.

Answer 'd' is incorrect as z should be the homogeneous function of x and y not non-homogeneous functions.

**Question 23**

If  $f(x, y) = \frac{x+y}{y}$ ,  $x \frac{dz}{dx} + y \frac{dz}{dy} = ?$

**(a) 0**

**(b) 1**

**(c) 2**

**(d) 3**

**Answer: a**

**Explanation:**

Given function  $f(x, y) = \frac{x+y}{y}$  can be written as  $f(x, y) = \frac{[1+\frac{y}{x}]}{\frac{y}{x}} = x^0 f(\frac{y}{x})$ ,

Hence by Euler's theorem.

$$x \frac{dz}{dx} + y \frac{dz}{dy} = 0$$

**Question 24**

**Find the approximate value of  $[0.982 + 2.012 + 1.942]^{1/2}$**

**(a) 1.96**

**(b) 2.96**

(c) 0.04

(d) -0.04

**Answer: b****Explanation:**Let  $f(x, y, z) = (x^2 + y^2 + z^2)^{\frac{1}{2}}$  .....(1)Hence,  $x = 1, y = 2, z = 2$ , so that  $dx = -0.02, dy = 0.01, dz = -0.06$ 

From (1)

$$\frac{df}{dx} = \frac{x}{f}$$

$$\frac{df}{dy} = \frac{y}{f}$$

$$\frac{df}{dz} = \frac{z}{f}$$

$$df = \frac{df}{dx}dx + \frac{df}{dy}dy + \frac{df}{dz}dz = \frac{(xdx + ydy + zdz)}{f} = \frac{-0.02 + 0.01 - 0.12}{3} = -0.04$$

$$[0.98^2 + 2.01^2 + 1.94^2]^{1/2} = f(1, 2, 2) + df = 3 - 0.04 = 2.96$$

**Question 25**

$f(x, y) = \frac{x^3 + y^3}{x^{99} + y^{98}x + y^{99}}$  find the value of  $f_y$  at  $(x, y) = (0, 1)$

(a) 101

(b) -96

(c) 210

(d) 0

**Answer: b****Explanation:**

Using Euler theorem

$$Xf_x + yf_y = n f(x, y)$$

Substituting  $x = 0; n = -96$  and  $y = 1$  we have

$$F_y = -96. F(0, 1) = -96.(1 / 1)$$

$$= -96$$

**Question 26**

$f(x, y) = x^3 + xy^2 + 901$  satisfies the Eulers theorem

(a) True

(b) False

(c) Not sure

(d) Invalid questions

**Answer: b****Explanation:**

The function is not homogenous and hence does not satisfy the condition posed by Euler's theorem.

**Question 27**

**For a homogenous function if critical points exist the value at critical points is**

- (a) 1 (b) equal to its degree  
(c) 0 (d) -1

**Answer: c**

For a homogeneous function if critical points exist the value at critical points is?  $f(a, b) = 0(a, b) \rightarrow$  critical points.  $nf(a, b) = 0 \Rightarrow f(a, b) = 0(a, b) \rightarrow$  critical points. Explanation: Euler's theorem is nothing but the linear combination asked here, The degree of the homogeneous function can be a real number.

**Question 28**

**$\lim_{n \rightarrow \infty} \left[ \frac{n}{1+n^2} + \frac{n}{4+n^2} + \frac{n}{9+n^2} + \dots + \frac{1}{2n} \right]$  is equal to**

- (a)  $\frac{\pi}{2}$  (b)  $\frac{\pi}{4}$   
(c) 1 (d) None of these

**Answer: d**

**Explanation:**

We have  $\lim_{n \rightarrow \infty} \left[ \frac{n}{1+n^2} + \frac{n}{4+n^2} + \dots + \frac{1}{2n} \right]$

$$= \lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{n}{r^2+n^2} = \lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{n}{n^2 \left(1 + \frac{r^2}{n^2}\right)}$$

$$= \lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{1}{n \left(1 + \frac{r^2}{n^2}\right)} = \int_0^1 \frac{dx}{1+x^2}$$

$$\left\{ \text{Applying formula, } \lim_{n \rightarrow \infty} \sum_{r=0}^{n-1} \left\{ f\left(\frac{r}{n}\right) \right\} \cdot \frac{1}{n} = \int_0^1 f(x) dx \right\}$$

$$= [\tan^{-1}x]_0^1 = \tan^{-1} 1 - \tan^{-1} 0 = \frac{\pi}{4}$$

**Question 29**

**For homogenous function with no saddle points we must have the minimum value as**

- (a) 90 (b) 1



(c) Equal to degree

(d) 0

**Answer: d****Explanation:**

Substituting  $f_x = f_y = 0$  At critical in euler's theorem we have  
 $nf(a, b) = 0 \rightarrow f(a, b) = 0(a, b) \rightarrow$  critical points.

**Question 30**

The derivates of  $f(x) = \int_{x^2}^{x^3} \frac{1}{\log t} dt$ , ( $x > 0$ ) is

(a)  $\frac{1}{3 \log x} - \frac{1}{2 \log x}$

(b)  $\frac{1}{3 \log x}$

(c)  $\frac{3x^2}{3 \log x}$

(d)  $(\log x)^{-1} \cdot x(x-1)$

**Answer: d****Explanation:****We know that**

$$\frac{d}{dx} \left( \int_a^b f(t) dt \right) = \frac{db}{dx} f(b) - \frac{da}{dx} f(a) \quad a \text{ and } b$$

Are functions of  $x \quad \therefore f(x) = \int_{x^2}^{x^3} \frac{1}{\log t} dt \quad e$

$$F'(x) = \frac{d}{dx}(x^3) \frac{1}{\log x^3} - \frac{d}{dx}(x^2) \frac{1}{\log x^2}$$

$$= \frac{3x^2}{3 \log x} - \frac{2x}{2 \log x} = x(x-1)(\log x)^{-1}$$

**Question 31**

The greatest value of the function  $f(x) = \int_1^x |t| dt$  on the interval  $\left[-\frac{1}{2}, \frac{1}{2}\right]$  is given by ?

(a)  $\frac{3}{8}$

(b)  $-\frac{1}{2}$

(c)  $-\frac{3}{8}$

(d)  $\frac{2}{5}$

**Answer: c****Explanation:**

$$f'(x) = |x| > 0 \forall x \in \left[-\frac{1}{2}, \frac{1}{2}\right] \text{ hence the}$$

Function is increasing on  $\left[-\frac{1}{2}, \frac{1}{2}\right]$  and therefore  $f(x)$  has  
Maximum at the right point of  $\left[-\frac{1}{2}, \frac{1}{2}\right]$

$$\text{Max } f(x) = f\left(\frac{1}{2}\right) = \int_1^{\frac{1}{2}} |t| dt = -\frac{3}{8}$$

### Question 32

**For homogenous function the linear combination of rates of independent change along x and y axis is**

- (a) Integral multiple function value      (b) no relation to function value  
(c) real multiple of function value      (d) depends if the function is a polynomial

**Answer: c**

**Explanation:**

Euler's theorem is nothing but the linear combination asked here, The degree of the homogeneous function can be a real number. Hence, the value is integral multiple of real number.

### Question 33

$\int_0^{b-c} f^n(x+a) dx$  = Homogenous function can be a real number. Hence the value is integral multiple of real number.

- (a)  $f'(a) - f'(b)$       (b)  $f'(b-c+a) - f'(a)$   
(c)  $f'(b+c-a) + f'(a)$       (d) None of these

**Answer: b**

**Explanation:**

$$\int_0^{b-c} f^n(x+a) dx$$

$$= [f'(x+a)]_0^{b-c} = f'(b-c+a) - f'(a).$$

### Question 34

$$\int_0^x \frac{x^3 dx}{(x^2+4)^2} =$$

- (a) 0      (b)  $\infty$   
(c)  $1/2$       (d) None of these

**Answer: b**

**Explanation:**

$$\begin{aligned} \int_0^{\infty} \frac{x^3 dx}{(x^2+4)^2} &= \frac{1}{2} \int_0^{\infty} \frac{2x^2 dx}{(x^2+4)^2} dx \\ &= 2 \int_0^{\infty} \frac{t}{(t+4)^2} dt, && \text{[Putting } x^2=t\text{]} \\ &= 2 \int_0^{\infty} \left[ \frac{1}{t+4} - \frac{4}{(t+4)^2} \right] dt = \frac{1}{2} \left[ \log(t+4) + \frac{4}{t+4} \right]_0^{\infty} \\ &= \frac{1}{2} [\log \infty + 0 - (\log 4 + 1)] = \infty \end{aligned}$$

**Question 35**

The points of intersection of  $F_1(x) = \int_2^x (2t - 5) dt$  and  $f_2(x) = \int_0^x 2t dt$ , are

- (a)  $\left(\frac{6}{5}, \frac{36}{25}\right)$  (b)  $\left(\frac{2}{3}, \frac{4}{5}\right)$   
 (c)  $\left(\frac{1}{3}, \frac{3}{6}\right)$  (d)  $\left(\frac{5}{4}, \frac{5}{7}\right)$

**Answer: a****Explanation:**

Let  $f_1(x) = y_1 = \int_2^x (2t - 5) dt$  and

$F_2(x) = y_2 = \int_0^x 2t dt$  now point of intersection means whose those point at which

$x^2 = x^2 - 5x + 6 \Rightarrow x = \frac{6}{5}$  and  $y = x^2 = \frac{36}{25}$  thus point of

Intersection is  $\left(\frac{6}{5}, \frac{36}{25}\right)$

**Question 36**

The solution of the equation  $\frac{x^2 d^2 y}{dx^2} = \ln x$ , when  $x=1, y=0$  and  $\frac{dy}{dx} = -1$

- (a)  $\frac{1}{2}(\ln x)^2 + \ln x$  (b)  $\frac{1}{2}(\ln x)^2 - \ln x$   
 (c)  $-\frac{1}{2}(\ln x)^2 + \ln x$  (d)  $-\frac{1}{2}(\ln x)^2 - \ln x$

**Answer: d****Explanation:**

$$\frac{d^2 y}{dx^2} = \frac{\log x}{x^2} \rightarrow \frac{-(\log x + 1)}{x} + C$$

$$\text{At } \frac{dy}{dx} = -\int \frac{\log x + 1}{x} dx = -\frac{1}{2}(\log x)^2 - \log x$$

**Question 37**

The rate of increase of bacteria in a certain culture is proportional to the number present. If it double 5 hours then in 25 hours its number would be

- (a) 8 times the original (b) 16 times the original  
(c) 32 times the original (d) 64 times the original

**Answer: c**

**Explanation:**

Let  $P_0$  be the initial population and let the

Population after  $t$  years be  $P$ . then  $\frac{dp}{dt} = KP \rightarrow \frac{dP}{P} = kdt$

On integrating, we have  $\log P = kt + c$  At  $t = 0$ ,

$P = P_0 \therefore \log P_0 = 0 + C, \therefore \log P = KT + \log P_0$

$\log \frac{P}{P_0} = kt$  when  $t = 5$  hrs,  $P = 2P_0 \therefore$

$\log \frac{2P}{P_0} = 5K \therefore K = \frac{\log 2}{5} \therefore \log \frac{P}{P_0} = \frac{\log 2}{5}t$  when

$T = 25$  hours, we have

$\log \frac{P}{P_0} = \frac{\log 2}{5} \times 25 = 5 \log 2 = \log 32; \therefore P = 32P_0.$

**Question 38**

The degree of the  $3\frac{d^2y}{dx^2} = \left\{1 + \left(\frac{dy}{dx}\right)^2\right\}^{\frac{3}{2}}$  is differential equation

- (a) 1 (b) 2  
(c) 3 (d) 6

**Answer: b**

**Explanation:**

$3\frac{d^2y}{dx^2} = \left\{1 + \left(\frac{dy}{dx}\right)^2\right\}^{\frac{3}{2}}$  on squaring, we

Get  $9\left(\frac{d^2y}{dx^2}\right)^2 = \left\{1 + \left(\frac{dy}{dx}\right)^2\right\}^3$  obviously the

Highest derivatives  $\frac{d^2y}{dx^2}$  contains a degree 2.

**Question 39**

The differential equation representing the family of curves  $y^2 = 2c(x + \sqrt{c})$ , where  $c$  is a positive parameter, is of

- (a) Order 1 (b) Order 2

(c) Degree 3

(d) Degree 4

**Answer: a****Explanation:**

Given family of curves

$$y^2 = 2c(x + \sqrt{c}), \quad \dots (i)$$

On differentiating both sides, we get

$$2y \frac{dy}{dx} = 2c(1 + 0) \rightarrow c = y \frac{dy}{dx}$$

From equation (i), we have

$$y^2 = 2y \frac{dy}{dx} \left\{ x + \left( y \frac{dy}{dx} \right)^{1/2} \right\}$$

$$\rightarrow \left( y^2 - 2xy \frac{dy}{dx} \right) = 2$$

**Question 41**

The order and degree of the differentiate equations  $\left(1 + 3 \frac{dy}{dx}\right)^{\frac{2}{3}} - 4$  are

(a)  $1, \frac{2}{3}$ 

(b) 3, 1

(c) 3, 3

(d) 1, 2

**Answer: c****Explanation:**

To check, order and degree, the given differential equation should be free from radicals, hence taking cube on both sides,

$$\left(1 + 3 \cdot \frac{dy}{dx}\right)^2 = \left(4 \cdot \frac{d^3y}{dx^3}\right)^3$$

Order = 3, degree = 3.

**Question 42**

The solution of the differential equation  $y - x \frac{dy}{dx} = a \left( y^2 + \frac{dy}{dx} \right)$  is

(a)  $y = c(x+a) (1+ay)$ (b)  $y = c(x+a) (1 - ay)$ (c)  $y = c(x-a) (1+ay)$ 

(d) None of these

**Answer: b****Explanation:**

$$Y - x \frac{dy}{dx} = a \left( y^2 + \frac{dy}{dx} \right)$$

$$Y - ay^2 = (x + a) \frac{dy}{dx} \quad \frac{dy}{y(1-ay)} = \frac{dx}{x+a}$$

On integrating both sides, we get  $\rho$

$$\text{Log } y - \log(1 - ay) = \log(x + a) + \log c$$

$$\frac{y}{(1-ay)} = c(x + a) \text{ or } c(x + a)(1 - ay) = y.$$

### **Question 43**

**Compute the sum of 4 digit numbers which can be formed with four digit 1, 3, 5, 7 if each digit is used once in each engagement:**

(a) 106646

(b) 106636

(c) 106666

(d) None of these

**Answer: d**

**Explanation:**

The number of arrangements of 4 different digits taken 4 at a time is given by  ${}^4P_4 = 4! = 24$ . All the four digits will occur equal number of times at each of the positions, namely ones, tens, hundreds, thousands.

Thus, each digit will occur  $\frac{24}{4} = 6$  times in each of the positions. The sum of digits in one's position will be  $6 \times (1+3+5+7) = 96$ . Similar is the case in ten's, hundred's and thousand's places.

Therefore, the sum will be  $96 + 96 \times 10 + 96 \times 100 = 106656$

### **Question 44**

**Eight guests have to be seated 4 on each side of a long rectangular table. 2 particular guests desire to sit on one side of the table and 3 on the other side. The number of ways in which the sitting arrangements can be made is:**

(a) 1732

(b) 1728

(c) 1730

(d) 1278

**Answer: b**

**Explanation:**

Let the two particular guests sit on right side.

So the three particular guests will sit on left side.

So remaining will be 3 people which need to be selected.

From these 3 people 2 will sit on right side and the one will sit on left side.

Total ways of arranging the people will be  $= {}^3C_2 \times {}^1C_1 = 3$

Total ways of arranging the people will be =  
 Selection of remaining  $\times 4!$  (For arranging people on left side)  $\times 4!$  (Arranging people on right side) =  $3 \times 24 \times 24 = 3 \times 756 = 1728$   
 So in 1728 ways we can arrange them

## PAST EXAMINATION QUESTIONS:

### MAY 2018

#### Question 1

The value of  $\int_1^2 \frac{1-x}{1+x} dx$  is equal to:

- (a)  $\log_2^3 - 1$  (b)  $2\log_2^3 - 1$   
 (c)  $\frac{1}{2}\log_2^3$  (d)  $\frac{1}{2}\log_2^3 - 1$

**Answer: b**

**Explanation:**

$$\begin{aligned} \int_1^2 \left( \frac{1-x}{1+x} \right) dx &= \int_1^2 \left( \frac{1}{1+x} - \frac{x}{1+x} \right) dx \\ &= \int_1^2 \frac{1}{1+x} dx - \int_1^2 \frac{x}{x+1} dx \\ &= \int_1^2 \frac{1}{1+x} dx - \int_1^2 \left( \frac{1+x-1}{1+x} \right) dx \\ &= \int_1^2 \frac{1}{(1+x)} dx - \int_1^2 \left( \frac{1}{1+x} \right) dx \\ &= \int_1^2 \frac{1}{1+x} dx - \int_1^2 1 \times dx + \int_1^2 \frac{1}{1+x} dx \\ &= 2 \int_1^2 \frac{1}{1+x} - \int_1^2 1 dx \\ &= 2[\log(1+x)]_1^2 - [x]_1^2 \\ &= 2[\log(2+1) - \log(1+1)] - [2-1] \\ &= 2[\log 3 - \log 2] - 1 \\ &= 2 \log \frac{3}{2} - 1 \end{aligned}$$

#### Question 2

$\int_0^2 \frac{3\sqrt{x}}{\sqrt{x}}$  is equal to

- (a)  $\frac{2\sqrt{2}}{\log e^3}$  (b) 0

(c)  $\frac{2(3\sqrt{2-1})}{\log_e 3}$

(d)  $\frac{3\sqrt{2}}{\sqrt{2}}$

**Answer: c****Explanation:**

$$\int_0^2 \frac{3^{\sqrt{x}}}{\sqrt{x}} dx$$

Let  $\sqrt{x} = t$

$$\int_0^2 3^{\sqrt{x}} \cdot \frac{1}{\sqrt{x}} dx \cdot \frac{1}{2\sqrt{x}} dx = dt$$

$$\frac{1}{\sqrt{x}} dx = 2dt$$

x	0	2
t	0	$\sqrt{2}$

$$\int_0^{\sqrt{2}} 3^t \cdot 2dt$$

$$\int_0^{\sqrt{2}} 3^t dt$$

$$2 \left[ \frac{3^t}{\log 3} \right]_0^{\sqrt{2}}$$

$$2 \left[ \frac{3^{\sqrt{2}}}{\log 3} - \frac{3^0}{\log 3} \right]$$

$$\frac{2(3^{\sqrt{2}} - 3^0)}{\log_e 3}$$

**Question 3****The value of  $\int_0^2 \frac{\sqrt{x}}{\sqrt{x} + \sqrt{2-x}} dx$  is:**

(a) 0

(b) 3

(c) 2

(d) 1

**Answer: d****Explanation:**

$$\int_0^2 \frac{\sqrt{x}}{\sqrt{x} + \sqrt{2-x}} dx \dots\dots\dots(1)$$

$$I = \int_0^2 \frac{\sqrt{0+2-x}}{\sqrt{0+2-x} + \sqrt{2-(0+2-x)}} dx$$

$$\left[ \int_a^b f(x) dx = \int_a^b f(a+b-x) dx \right]$$

$$I = \int_0^2 \frac{\sqrt{2-x}}{\sqrt{2-x} + \sqrt{x}} dx \dots\dots\dots(2)$$

**Apply (1) and (2) we get**



$$2 I = \int_0^2 \left[ \frac{\sqrt{x}}{\sqrt{x+\sqrt{2-x}}} + \frac{\sqrt{2-x}}{\sqrt{2-x+\sqrt{x}}} \right] dx$$

$$2 I = \int_0^2 \frac{(\sqrt{x} + \sqrt{2-x})}{(\sqrt{x} + \sqrt{2-x})} dx$$

$$2 I = \int_0^2 1 dx$$

$$2 I = [X]_0^2$$

$$2 I = [2 - 0]$$

$$2 I = 2 - 0 = 2$$

#### Question 4

$$\lim_{X \rightarrow 1} \frac{x + x^2 + x^3 \dots \dots + x^n - n}{x - 1}$$

(a) n

(b)  $\frac{n(n+1)}{2}$

(c) (n + 1)

(d) n(n + 1)

**Answer: b**

**Explanation:**

$$\lim_{X \rightarrow 1} \frac{x + x^2 + x^3 \dots \dots + x^n - n}{x - 1} \quad (:)$$

**By L.H. Rule**

$$\begin{aligned} &= \lim_{x \rightarrow 1} \frac{d/dx(x+x^2+x^3+\dots+x^n-n)}{d/dx(x-1)} \\ &= \lim_{x \rightarrow 1} \frac{1+2x+3x^2+\dots+nx^{n-1}-0}{1-0} \\ &= \frac{1+2 \times 1+3(1)^2+\dots+n(1)^{n-1}}{1} \\ &= 1 + 2 + 3 + \dots + n \\ &= \sum_{n=1}^n \frac{n(n+1)}{2} \end{aligned}$$

#### Question 5

The cost function for the production of x unit of a commodity is given by  $C(x) = 2x^3 + 5x^2 + 36x + 15$

(a) 3

(b) 2

(c) 1

(d) 4

**Answer: a**

**Explanation:**

The cost function given by  $C(x) = 2x^3 + 15x^2 + 36x + 15$

$$\frac{d}{dx} C(x) = 6x^2 - 30x + 36 \dots\dots(1)$$

$$C(x) = 0$$

$$6x^2 - 30x + 36 = 0$$

$$6(x^2 - 5x + 6) = 0$$

$$= x^2 - 5x + 6 = 0$$

$$= x^2 - 3x - 2x + 6 = 0$$

$$= x(x-3) - 2(x-3) = 0$$

$$(x-3)(x-2) = 0$$

$$X = 3, 2$$

Differentiating equations (2) again w.r.f. 'x'

$$C(x) = 12x - 30 \quad \text{Eq} \dots\dots\dots (3)$$

Putting (x = 2) in

$$C(x) = 12 \times 2 - 30 = -6$$

Putting (x=3) in

$$C(x) = 12 \times 3 - 30 = 6(+ve) \text{ so function is minimum at } x=3$$

### **Question 6**

$$\lim_{x \rightarrow 0} \frac{2e^{\frac{1}{x}-3x}}{e^{\frac{1}{x}+x}}$$

(a) -3

(b) 0

(c) 2

(d) 9

### **Answer: C**

$$\text{Let } \frac{1}{x} = y \text{ if } x \rightarrow 0, y \rightarrow \infty$$

$$\lim_{x \rightarrow \infty} \frac{2e^y - 3\frac{1}{y}}{e^y + \frac{1}{y}}$$

$$= \lim_{x \rightarrow \infty}$$

$$\frac{2 - 3\frac{1}{\infty \cdot e^\infty}}{1 + \frac{1}{\infty \cdot e^\infty}}$$

$$= \frac{2 - 0}{1 + 0} = 2$$

$$= \frac{2-0}{1+0} = 2$$

**NOV 2018****Question 1**

Let  $x = at^3$ ,  $y = \frac{a}{t^2}$ . Then  $\frac{dy}{dx} =$

(a)  $\frac{-1}{t^6}$

(b)  $\frac{-3a}{t^6}$

(c)  $\frac{1}{3at^6}$

(d) None

**Answer: d****Explanation:**

$$\text{If } x = at^3, y = \frac{a}{t^2} = at^{-2}$$

$$\text{Given } x = at^3$$

Different w.r.t. (t)

$$\frac{dy}{dx} = \frac{d}{dt} at^3 = a \cdot 3t^2 = 3at^2$$

and  $y = at^{-2}$ 

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{-2at^{-3}}{3at^2} = \frac{-2}{3t^5}$$

**Question 2**

$\int x(x^2 + 4)^5 dx$  is equal to

(a)  $(x^2 + 4)^6 + c$

(b)  $\frac{1}{12}(x^2 + 4)^6 + c$

(c)  $\frac{1}{6}(x^2 + 4)^6 + c$

(d) None

**Answer: b****Explanation:**

$$\int x(x^2 + 4)^5 dx = x$$

$$\text{Let } x^2 + 4 = t$$

$$2x dx = dt$$

$$x dx = \frac{dt}{2}$$

$$\int (x^2 + 4)^5 \cdot x dx$$

$$\int t^5 = \frac{dt}{2}$$

$$= \frac{1}{2} \int t^5 dt$$

$$= \frac{1}{2} \frac{t^6}{6} + c$$

$$= \frac{1}{12} (x^2 + 4)^6 + c$$

**Question 3**

$$xy = 1 \text{ then } y^2 + \frac{dy}{dx} = ?$$

- (a) 1 (b) 0  
(c) 2 (d) None

**Answer: b****Explanation:**

$$Xy = 1 \text{ then } y^2 + \frac{dy}{dx} = ?$$

$$\text{Given } xy = 1$$

$$Y = \frac{1}{x} \text{----- (1)}$$

$$Y = x^{-1}$$

$$\frac{dy}{dx} = (-1)x^{-2}$$

$$\frac{dy}{dx} = \frac{-1}{x^2}$$

**Question 4**

$$\int_{-1}^3 (1 + 3x + x^3) dx \text{ is equal to}$$

- (a) -4 (b) 4  
(c) 3 (d) -3

**Answer: a**

$$\int_{-1}^3 (1 + 3x + x^3) dx$$

$$\int_{-1}^3 1 dx + \int_{-1}^3 3x dx - \int_{-1}^3 x^3 dx$$

$$[x]_{-1}^3 + 63 \left[ \frac{x^2}{2} \right]_{-1}^3 - \left[ \frac{x^4}{4} \right]_{-1}^3$$

$$[3 - (-1)] + \frac{3}{2} [(3)^2 - (-1)^2] - \frac{1}{4} [(3)^4 - (-1)^4]$$

$$(3+1)+\frac{3}{2}[9-1]-\frac{1}{4}[81-1]$$

$$4+\frac{3}{2}\times 8-\frac{1}{4}\times 80$$

$$4+12-20 = -4$$

## MAY 2019

### Question 1

If  $2^x - 2^y = 2^{x-y}$  then  $\frac{dy}{dx}$  at  $x = y = 2$

- (a) 1 (b) 2  
(c) 4 (d) 5

**Answer: a**

**Explanation:**

$$2^x - 2^y = 2^{x-y} \quad x = y = 2 \quad \frac{dy}{dx}$$

$$2^x \cdot \text{Log}^2 - 2^y \cdot \log^2 \cdot \frac{dy}{dx} = 2^{x-y} \cdot \text{Log}^2 \left[ 1 - \frac{dy}{dx} \right]$$

$$\text{Log}^2 [2^x - 2^y \cdot \frac{dy}{dx}] = \text{Log}^2 [2^{x-y} (1 - \frac{dy}{dx})]$$

$$2^2 - 2^2 \cdot \frac{dy}{dx} = 2^0 \left[ 1 - \frac{dy}{dx} \right]$$

$$4 - 4 \cdot \frac{dy}{dx} = 1 - \frac{dy}{dx}$$

$$4 - 1 = 4 \frac{dy}{dx} - \frac{dy}{dx}$$

$$3 = 3 \frac{dy}{dx}$$

$$\frac{dy}{dx} = 1$$

### Question 2

If the cost of function of a commodity is given by  $C = 150x - 5x^2 + \frac{x^3}{6}$ , where C stands for cost and x stands for output. If the average cost is equal to the marginal cost then the output x = \_\_\_\_\_

- (a) 5 (b) 10  
(c) 15 (d) 20

**Answer: c****Explanation:**

$$\text{Average cost} = \frac{\text{Total cost}}{\text{output}}$$

$$C = 150x - 5x^2 + \frac{x^3}{6}$$

$$\frac{C}{\text{output}} = \frac{150x}{x} - \frac{5x^2}{x} + \frac{x^3}{6x}$$

$$C = 150 - 5x + \frac{x^2}{6}$$

$$\frac{dC}{dx} = -5 + \frac{2x}{6}$$

$$-5 + \frac{x}{3} = 0$$

$$75 + x = 0$$

$$X = 15$$

**Question 3**

$$\int_2^3 \frac{\sqrt{x}}{\sqrt{5-x} + \sqrt{x}} dx =$$

(a) 1

(b)  $\frac{1}{2}$

(c) 2

(d)  $\frac{3}{2}$

**Answer: a****Explanation:**

$$I = \int_2^3 \frac{\sqrt{x}}{\sqrt{5-x} + \sqrt{x}} \dots\dots(1)$$

$$I = \int \frac{\sqrt{5-x}}{\sqrt{15-5+x} + \sqrt{5-x}} = \int \frac{\sqrt{5-x}}{\sqrt{x} + \sqrt{5-x}} \dots\dots(2)$$

$$2I = \int_2^3 \frac{\sqrt{x}}{\sqrt{5-x} + \sqrt{x}} + \int \frac{\sqrt{x}}{\sqrt{5-x} + \sqrt{x}}$$

$$2I = \frac{\sqrt{5} + \sqrt{5-x}}{\sqrt{5-x} + \sqrt{x}}$$

$$2I = 2$$

$$X = 1$$

**Question 4**

$$\int \log_e(a^x) dx =$$

(a)  $\log_e a \left[ \frac{x^2}{2} \right] + c$

(b)  $\log_e a \left[ \frac{x}{2} \right] + c$

(c)  $x \log_e a^x - x + c$

(d) None of these

**Answer: a****Explanation:**

$$\int \log_e (a^x) dx$$

By option method: Base method

Differentiate option a

$$\log_e a^{\left[\frac{x^2}{2}\right]}$$

$$\frac{1}{a^{\left[\frac{x^2}{2}\right]}} \times a^{\left[\frac{x^2}{2}\right]} \cdot \log a \cdot x \cdot \frac{2x}{2}$$

$$= x \cdot \log_e a^x$$

$$= \log \frac{a^x}{e}$$

## NOV 2019

### Question 1

$$\int a^x dx.$$

(a)  $x^x(1 + \log x)$

(c)  $x \cdot \log x$

(b)  $1 + \log x$

(d)  $\frac{a^x}{\log a} + c$

**Answer: d****Explanation:**

(d) Since, we know that

$$\frac{d}{dx} \frac{a^x}{\log a} = a^x$$

$$\int a^x dx = \frac{a^x}{\log a} + c$$

### Question 2

$$\int x \cdot e^x dx.$$

(a)  $e^x (1 + \log x)$

(c)  $\log x + e^x + c$

(b)  $e^x \cdot x + e^x + c$

(d)  $\frac{x^2}{e^x} + c$

**Answer: a**

**Explanation:**

(a)  $\int x \cdot e^x dx$ .

Following I = Inverse

L = Logarithmic

A = Algebraic

T = Trigonometric

E = Exponential

So,  $x \Rightarrow$  I<sup>st</sup> function $e \Rightarrow$  II<sup>nd</sup> function $x \cdot e^x \quad x \Rightarrow u$  $\int I \cdot II^{dx} \quad \text{or} \quad e^x \Rightarrow v$ 

Property

Since,  $\int u \cdot v dx = u \cdot \int v dx - \int \left[ \frac{d}{dx}(u) \cdot \int v dx \right] dx$

$x \cdot \int e^x dx - \int \left[ \frac{d}{dx}(x) \cdot \int e^x dx \right] dx$

$x \cdot e^x - \int [1 \times e^x] dx$

$x \cdot e^x - e^x + c$

$e^x (x - 1) + c$

**Question 3**

$\int (4x + 3)^6 dx$ .

(a)  $\frac{1}{28} (4x + 3)^7 + c$

(b)  $\frac{1}{7} (4x + 3)^7 + c$

(c)  $\frac{1}{6} (4x + 3)^6 + c$

(d)  $\frac{4x}{5} + \frac{3}{5} + c$

**Answer: a****Explanation:**

(a)  $\int (4x + 3)^6 dx$

Since,  $\int x^n dx = \frac{x^{n+1}}{n+1} + c$

$\int (ax + b)^n dx = \frac{(ax+b)^{n+1}}{(n+1)a} + c$

So,

$\int (4x + 3)^6 dx$

$= \frac{(4x+3)^{6+1}}{(6+1) \cdot 4} + c$

$= \frac{1}{28} (4x + 3)^7 + c$



**Question 4**

$$\int_{-1}^1 (2x^2 - x^3) dx$$

(a)  $\frac{4}{3}$

(b) 1

(c) 2

(d)  $\frac{2}{3}$

**Answer: a****Explanation:**

(a)  $\int_{-1}^1 (2x^2 - x^3) dx$

$$= \left[ 2 \times \frac{x^3}{3} - \frac{x^4}{4} \right]_{-1}^1$$

$$= \left[ \left( \frac{2}{3} \times 1^3 - \frac{1^4}{4} \right) - \left\{ \frac{2}{3} \times (-1)^3 - \frac{(-1)^4}{4} \right\} \right]$$

$$= \left[ \left( \frac{2}{3} - \frac{1}{4} \right) - \left( -\frac{2}{3} - \frac{1}{4} \right) \right]$$

$$= \frac{2}{3} - \frac{1}{4} + \frac{2}{3} + \frac{1}{4}$$

$$= \frac{4}{3}$$

**Question 5**

$\frac{d}{dx}(x \cdot \log x)$

(a)  $x(1 + \log x)$

(b)  $1 + \log x$

(c)  $e^x x \cdot \log x$

(d)  $x^2 (\log x)$

**Answer: b****Explanation:**

(b)  $\frac{d}{dx}(x \cdot \log x)$

$$\text{Since } \frac{d}{dx}(u \cdot v) = u \frac{d}{dx}(v) + v \frac{d}{dx}(u)$$

$$\text{So here } u \Rightarrow x$$

$$v \Rightarrow \log x$$

$$\therefore \frac{d}{dx}(x \cdot \log x)$$

$$= x \cdot \frac{d}{dx}(\log x) + \log x \times \frac{d}{dx}(x)$$

$$= x \cdot \frac{1}{x} + \log x \times 1$$

$$= 1 + \log x$$

**Question 6**

**Differentiate  $x^x$  w.r.t  $x$ .**

(a)  $x^x (1 + \log x)$

(b)  $\frac{y}{x}$

(c)  $\frac{-y}{x}$

(d)  $y + x^x \log x$

**Answer: a****Explanation:**

(a)  $\frac{d}{dx} (xx) = ?$

Net  $y = xx$

Using log both sides

$\log y = x \log x$

On differentiating both sides w.e.t.  $x$ 

$$\frac{1}{y} \cdot \frac{dy}{dx} = x \times \frac{d}{dx} (\log x) + \log x \times \frac{d}{dx} (x)$$

$$\frac{dy}{dx} = y \left[ x \times \frac{1}{x} + \log x \times 1 \right]$$

**Question 7**

$$\int x^2 \cdot e^x dx.$$

(a)  $2x \cdot e^x$

(b)  $e^x (x^2 - 2x)$

(c)  $x^2 \cdot e^x \cdot (2x) + 2$

(d)  $e^x (x - 1)$

**Answer: C****Explanation:**

$$\int x^2 e^x dx$$

Using I late

$a^2 \Rightarrow 1^{\text{st}} \text{ function (u)}$

$e^x \Rightarrow 2^{\text{nd}} \text{ function (v)}$

$$\int u \cdot v dx = u \cdot \int \left[ \frac{d}{dx} (u) \cdot \int v dx \right] dx$$

So  $\int x^2 e^x dx$

$$x^2 \int e^x dx - \int \left[ \frac{d}{dx} (x^2) \int e^x dx \right] dx$$

$$x^2 e^x dx - \int [2x \cdot e^x] dx$$

$$x^2 \cdot e^x - 2x \int x \cdot e^x dx \quad \text{-----Equation (1)}$$

$$= x \cdot \int e^x - \int \frac{d}{dx} (x) \cdot \int e^x dx dx$$

$$= x \cdot e^x - e^x$$

$$= e^x (x - 1) \quad \text{-----Equation (2)}$$

Put Equation (2) in Equation (1)

$$x^2 \cdot e^x - 2 e^x (x-1)$$

$$x^2 \cdot e^x - 2 e^x \cdot x + 2$$

## JULY 2021

### Question 1

The value of  $\int_{-2}^2 f(x) dx$ , where  $f(x) = 1+x, x \leq 0$ ;  $f(x) = 1-2x, x \geq 0$  is

- (a) 20  
(b) -2  
(c) -4  
(d) 0

**Answer: Options (b)**

## DEC 2021

### Question 1

The cost of producing  $x$  units is  $500-20x^2 + x^3 / 3$ . The marginal cost is minimum at  $x$  = \_\_\_\_\_.

- (a) 5  
(b) 10  
(c) 40  
(d) 50

**Answer: c**

**Explanation:**

Here, cost function is given by

$$c(x) = 500 - 20x^2 + \frac{x^3}{3}$$

Diff. w.r.t. 'x'

$$\frac{d}{dx} c(x) = \frac{d}{dx} \left[ 500 + 20x^2 + \frac{x^3}{3} \right]$$

$$\frac{dc(x)}{dx} = 0 - 40x + \frac{3x^2}{3}$$

$$\frac{dc}{dx} = (x^2 - 40x)$$

$$\text{Marginal cost} = \frac{dc}{dx}$$

$$= (x^2 - 40x)$$

$$x(x-40) = 0$$

$$\text{If } x=0, \text{ if } x-40 = 0$$

$$x = 40$$

### **Question 2**

If  $y = \frac{x^4}{e^x}$  then  $\frac{dy}{dx}$  is equal to:

(a)  $x^3(4-x) / (e^x)^2$

(b)  $x^3(4-x) / e^x$

(c)  $x^2(4-x) / e^x$

(d)  $x^3(4x-1) / e^x$

**Answer: b**

**Explanation:**

$$\text{If } y = \frac{x^4}{e^x}$$

Diff. w.r.t. 'x'

$$\begin{aligned} \frac{dy}{dx} &= \frac{e \times \frac{d}{dx}(x^4) - x^4 \cdot e^x}{(e^x)^2} \\ &= \left( \frac{e^x \cdot 4x^3 - x^4 \cdot e^x}{e^{2x}} \right) \\ &= \frac{x^3(4-x)}{e^x} \end{aligned}$$

### **Question 3**

The speed of a train at a distance  $x$  (from the starting point) is given by  $3x^2 - 5x + 4$ .

What is the rate of change (of distance) at  $x=1$ ?

(a) -1

(b) 0

(c) 1

(d) 2

**Answer: c**

**Explanation:**

The speed of a train at a distance  $x$  is given by

$$V = 3x^2 - 5x + 4$$

Diff. w.r.t. 'x'

$$\frac{dy}{dx} = 6x - 5$$

$$\left[ \frac{dy}{dx} \right]_{x=1} = 6 \times 1 - 5 = 6 - 5 = 1$$

Rate of change (of distance) at  $x = 1$  is 1.

**JUNE 2022****Question 1**

$\int_0^1 x e^x dx$  is equal to:

- (a) 0 (b) 2  
(c) 1 (d) 3

**Answer: Options (c)**

**Explanation:**

$$\int_0^1 x e^x dx$$

$$\left[ x \int e^x dx - \int \left( \frac{d}{dx} x \int e^x dx \right) dx \right]_0^1$$

$$= [x e^x - \int 1 \cdot e^x dx]_0^1$$

$$= [x e^x - e^x]_0^1$$

$$= (1 \cdot e^1 - e^1) - (0 \cdot e^0 - e^0)$$

$$= (e - e) - (0 - 1) = 0 + 1 = 1$$

**Question 2**

What will be  $f(x)$  if  $f'(x) = 10x^2 + 4x$  and  $f(-3) = 17$

- (a)  $f(x) = \frac{10x^3}{3} + 2x^2 + 89$  (b)  $f(x) = \frac{10x^3}{3} + 2x^2 + 72$   
(c)  $f(x) = \frac{10x^3}{3} + 2x^2 - 89$  (d) None

**Answer: Options (a)**

**Explanation:**

Here  $f'(x) = 10x^2 + 4x$

on integration both side

$$\int f'(x) dx = \int (10x^2 + 4x) dx$$

$$f(x) = 10 \frac{x^3}{3} + 4 \frac{x^2}{2} + C \quad (1)$$

putting  $x = -3$ ,  $f(-3) = \frac{10(-3)^3}{3} + \frac{4(-3)^2}{2} + C$

$$17 = \frac{10(-27)}{3} + \frac{4 \times 9}{2} + c$$

$$17 = -90 + 18 + c$$

$$c = 89$$

putting  $C = 89$  in eq (1)

$$f(x) = 10\frac{10x^3}{3} + \frac{4x^3}{2} + 89$$

$$f(x) = 10\frac{10x^3}{3} + 2x^2 + 89$$

### Question 3

$\int (\log x)^2 dx$  is equal to:

- (a)  $x(\log x)^2 - 2x \log x + 2x + C$   
 (c)  $x(\log x)^2 - 2x \log x - x + C$

- (b)  $x(\log x)^2 + 2x \log x - 2x + C$   
 (d) None

**Answer: Options (a)**

**Explanation:**

$$\begin{aligned} I &= \int (\log x)^2 dx \\ &= \int (\log x)^2 \cdot 1 dx \\ &= (\log x)^2 \cdot \int 1 dx - \int \left(\frac{d}{dx}(\log x)^2 \cdot \int 1 dx\right) dx \\ &= (\log)^2 \cdot x - \frac{2 \log x}{x} \cdot x dx \\ &= x(\log x)^2 - 2[\log x \int 1 dx - \int \left(\frac{d}{dx} \log x \cdot \int 1 dx\right) dx] \\ &= x(\log)^2 - 2[\log x \cdot (x) - \int \frac{1}{x} \cdot x dx] \\ &= x(\log x)^2 - 2[\log x \cdot (X) - \int \frac{1}{x} \cdot x dx] \\ &= x(\log)^2 - 2[x \log x - x] + C \\ &= x(\log)^2 - 2x \log x + 2x + C \end{aligned}$$

### Question 4

The derivative of the function  $\sqrt{x + \sqrt{x}}$  is

(a)  $\frac{1}{2\sqrt{x+\sqrt{x}}}$

(b)  $1 + \frac{1}{2\sqrt{x}}$

(c)  $\frac{1}{2\sqrt{x+\sqrt{x}}} \left(1 + \frac{1}{2\sqrt{x}}\right)$

(d) None of these

**Answer: Options (c)**

**Explanation:**

$$y = \sqrt{x + \sqrt{x}}$$

Diff w. r. t 'a'

$$\begin{aligned} \frac{dy}{dx} &= \frac{d}{dx} (\sqrt{x + \sqrt{x}}) \\ &= \frac{1}{2\sqrt{x+\sqrt{x}}} \left(1 + \frac{1}{2\sqrt{x}}\right) \end{aligned}$$